

Table A22 Notifications and Deaths due to Tuberculosis 1961 to 2000

Year	Notification Rate (Number of Cases per 100 000 Population)	Total Deaths from Tuberculosis	Tuberculosis Death Rate (Number of Deaths per 100 000 Population)	Average Age at Death from Tuberculosis
1961	397.21	1 907	60.19	43.0
1966	314.80	1 515	41.74	53.0
1971	223.17	1 250	30.90	57.5
1976	175.48	568	12.57	59.5
1981	149.11	489	9.43	63.0
1982	142.98	454	8.62	63.0
1983	136.59	446	8.34	63.0
1984	145.30	420	7.78	64.5
1985	138.28	409	7.50	65.5
1986	134.53	407	7.37	68.0
1987	130.26	405	7.26	68.5
1988	124.76	388	6.89	69.0
1989	117.90	403	7.09	69.0
1990	114.12	382	6.70	69.0
1991	109.23	409	7.11	69.0
1992	112.65	410	7.07	68.0
1993	110.78	396	6.71	69.0
1994	104.70	409	6.78	71.0
1995	100.91	418	6.79	71.1
1996	101.02 *	292	4.54 *	70.6
1997	108.98 *	252	3.88 *	72.1
1998	117.26 *	270	4.13 *	72.6
1999	113.71 *	312	4.72 *	72.9
2000	113.70 *	299	4.49 *	73.4

Notes Notification figures in the table refer to the tuberculosis cases known to the Department of Health.

\* Since August 2000, the "resident population" approach has been adopted in place of the "extended de facto" approach for compiling population estimates. Also, the Population Census which was conducted in March 2001 provides a benchmark for revising the population estimates compiled since the 1996 Population By-census. In this table, the population-related figures from 1996 onwards have been revised accordingly.