監測及流行病學處



Surveillance And Epidemiology Branch

保障市民健康 Protecting Hong Kong's health

本署檔號 Our Ref. : (105) in DH SEB CD/8/71/1

來函檔號 Your Ref : 電 話 Tel. : 傳 真 Fax No. :

2 March 2009

Dear Doctor,

<u>Updated Vaccination Schedule for Pneumococcal Conjugate Vaccine</u> <u>Under Childhood Immunisation Programme</u>

I am pleased to inform you that the Scientific Committee on Vaccine Preventable Diseases (SCVPD) of the Centre for Health Protection (CHP) has provided updated recommendations on the use of 7-valent pneumococcal conjugate vaccine (PCV7) under the Government's Childhood Immunisation Programme (CIP).

Starting from the 3rd quarter of 2009, PCV7 will be incorporated in CIP and the vaccination schedule as recommended by SCVPD will be adopted. Details of the recommendations are summarised as follows for your information-

Standard	• A standard 3-dose primary series at 2 nd , 4 th and 6 th months	
Regimen	of age with a bo	poster at 12-15 months
Catch-up	Children aged 6	• A 3-dose primary series at any time
Schedule	months or below	with 4-8 weeks' interval between
		doses; a booster dose at 12-15
		months or 2 months after the last
		dose whichever is later (i.e. 3+1).
	Children aged	• A 2-dose primary series with an
	between 7 months	interval of 4-8 weeks but not later
	and less than 1	than the age of 1 year; a booster dose
	year	at 12-15 months with an interval of
		at least 2 months after the last dose
		(i.e. 2+1)
		• If the second dose of the primary
		series is not administered by the age

	of 1 year, a booster dose at 12-15 months with an interval of at least 2 months after the last dose (i.e. 1+1).
Children aged	One single dose
between 1 year and	
less than 2 years	

You may also wish to visit CHP website (www.chp.gov.hk) for the complete SCVPD's recommendations on the use of pneumococcal vaccines.

Thank you for your attention.

Yours sincerely,

(Dr SK CHUANG)

Consultant Community Medicine
(Communicable Disease)
Centre for Health Protection